



BOVISTART

Complementary feed ready to use for newborn calves

PROPERTIES

BOVISTART provides to newborn calves the essential elements needed at birth:

- Energy supply in the form of highly digestible sugars and essential fatty acids
- Vitamins, essential for the vitality of newborn calves
- Natural lactic acid bacteria, promoting a proper development of the digestive lactic flora
- Colostrum to help to strengthen natural defences, for the good health of calves.

COMPOSITION

Dried egg products
Cod liver oil
Lactose
IBR free colostrum
Dextrose

ANALYTICAL CONSTITUENTS

Moisture	48.3 %
Crude protein	12.7 %
Crude fibre	0.3 %
Crude fats	20.6%
Crude ash (mineral matter)	1.7 %
Sodium	0.25 %

ADDITIVES (per liter) (Density: 1.05)

Gut flora stabilizer

4b1705i *Enterococcus lactis* NCIMB 10415: 1.0x10¹¹ CFU/L

Vitamins

3a672b Vitamin A	1 250 000 IU/L
3a671 Vitamin D3	440 000 IU/L
3a700 Vitamin E	940 IU/L
3a300 Vitamin C	6 450 mg/L
3a820 Vitamin B1	250 mg/L
3a825ii Vitamin B2	1 000 mg/L

Preservatives

1a330 Citric acid	10 000 mg/L
1k202 Potassium sorbate	5 000 mg/L

Antioxygens

E310 Propyl gallate	60 mg/L
1b320 Butylated hydroxyanisole (BHA)	30 mg/L

Aromatic substances

Mix of aromatic substances
2b139-eo Essential oil lemon exprimée : 2 550 mg/L

DIRECTIONS OF USE

BOVISTART is an oral paste ready and easy to use thanks to its syringe. It is orally distributed to the newborn calf:

- One syringe per calf in the first hours after birth
- For the weakest calves, who do not suckle, do not drink or had a difficult birth, repeat with a second syringe the day after
- One syringe of **BOVISTART** is also recommended for older calves who would lack of vigor, for a “boost” effect.

Precautions for use

- Before use or before extending the period of use, it is recommended to seek the advice of a feed expert or a veterinarian
- The simultaneous use of different organic acids or their salts is contraindicated where one or more of them is used at or near the maximum permitted content.

PACKAGING

- Blister of 6 syringes of 15 mL each
- Box of 25 syringes of 15 mL each
- Should be used within 12 months after manufacture
- **BOVISTART** must be stored in a temperature range from 10 to 20°C.

